

BLACK-BRANE SOLUTION FOR A_3 ALGEBRA

M.A. Grebeniuk¹, V.D. Ivashchuk² and V.N. Melnikov³

*Center for Gravitation and Fundamental Metrology, VNIIMS,
3/1 M. Ulyanovoy Str., Moscow 117313, Russia and*

*Institute of Gravitation and Cosmology, People's Friendship University
of Russia, Mikhlukho-Maklaya Str. 6, Moscow 117198, Russia*

Abstract

Black p -brane solutions for a wide class of intersection rules and Ricci-flat “internal” spaces are considered. They are defined up to moduli functions H_s obeying non-linear differential equations with certain boundary conditions imposed. A new solution with intersections corresponding to the Lie algebra A_3 is obtained. The functions H_1 , H_2 and H_3 for this solution are polynomials of degree 3, 4 and 3, correspondingly. An example of A_3 -solution with three 3-branes in 12-dimensional model (suggested by N. Khviengia et al) is presented.

PACS: 04.20.Jb; 04.50.+h; 04.70.Bw; 02.20.Sv; 02.30.Hq

Keywords: Black-branes; polynomials; Lie algebras

¹mag@gravi.phys.msu.su

²ivas@rgs.phys.msu.su

³melnikov@rgs.phys.msu.su

1 Introduction

In this paper we continue investigation of the black-brane solutions with arbitrary intersection rules [1, 2, 3, 4]. These solutions are governed by functions $H_s = H_s(R)$ obeying a set of second order non-linear differential equations (equivalent to Toda-type ones) with certain boundary conditions imposed.

More general spherically symmetric solutions were obtained in [14] using the sigma-model approach from [7, 13, 4]. For another (algebraic) approach see also [5, 6, 9] and references therein.

In order to simplify the investigation of such model and obtain exact solutions the following conjecture was suggested in [1]: the functions H_s are polynomials if the p -brane intersection rules correspond to semisimple Lie algebras. This conjecture was verified for solutions with A_m and C_m series of Lie algebras [2, 3] and also confirmed by special black-brane "block-orthogonal" solutions considered earlier in [15, 16]. It should be noted that an analogue of this conjecture for extremal black holes was considered earlier in [11].

Although the conjecture mentioned above allows one to reduce the set of differential equations to the system of algebraic one, it still seems to be very difficult to get the general solution to this set of equations too. So, the only way of obtaining exact solutions is to investigate special cases with different Lie algebras (such as A_2 , C_2 etc.).

The simplest nontrivial example of black-brane solution corresponding to Lie algebra $A_2 = sl(3)$ was obtained in [1, 2] (e.g. dyonic solutions in 11-dimensional supergravity). This A_2 -solution was governed by two polynomials H_1 and H_2 of degree 2. The coefficients of polynomials and charges were the functions of some parameters P_1 and P_2 and had a rather simple form.

Here we present a new solution corresponding to the Lie algebra A_3 and governed by three polynomials of degree 3, 4 and 3. The coefficients of polynomials exhibit a non-meromorphic dependence upon parameters P_1 , P_2 and P_3 : a term, which is a square root of a polynomial of degree 4, appears. (For the special case with $P_1 = P_3$ the solution was already obtained and investigated in [18]). Here we also suggest an example of A_3 -solution with three 3-branes in 12-dimensional model from [19].

2 p -brane black hole solutions

Consider a model governed by the action

$$S = \int d^D x \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2 \right\} \quad (1)$$

where $g = g_{MN}(x) dx^M \otimes dx^N$ is a metric, $\varphi = (\varphi^\alpha) \in \mathbb{R}^l$ is a vector of scalar fields, $(h_{\alpha\beta})$ is a constant symmetric non-degenerate $l \times l$ matrix ($l \in \mathbb{N}$), $\theta_a = \pm 1$,

$$F^a = dA^a = \frac{1}{n_a!} F_{M_1 \dots M_{n_a}}^a dz^{M_1} \wedge \dots \wedge dz^{M_{n_a}} \quad (2)$$

is a n_a -form ($n_a \geq 1$), λ_a is a 1-form on $\mathbb{R}^l$: $\lambda_a(\varphi) = \lambda_{\alpha a} \varphi^\alpha$, $a \in \Delta$, $\alpha = 1, \dots, l$. In (1) we denote $|g| = |\det(g_{MN})|$, $(F^a)_g^2 = F_{M_1 \dots M_{n_a}}^a F_{N_1 \dots N_{n_a}}^a g^{M_1 N_1} \dots g^{M_{n_a} N_{n_a}}$, $a \in \Delta$. Here Δ is some finite set. It should be noted that in the models with one time all $\theta_a = 1$ if the signature of the metric is $(-1, +1, \dots, +1)$.

Let us consider (black-brane) solutions to the field equations corresponding to the action (1) from [1, 2, 3]. These solutions are defined on the manifold

$$M = (R_0, +\infty) \times (M_1 = S^{d_1}) \times (M_2 = \mathbb{R}) \times \dots \times M_n, \quad (3)$$

and have the following form

$$g = \left(\prod_{s \in S} H_s^{2h_s d(I_s)/(D-2)} \right) \left\{ f^{-1} dR \otimes dR + R^2 d\Omega_{d_1}^2 \right. \quad (4)$$

$$\left. - \left(\prod_{s \in S} H_s^{-2h_s} \right) f dt \otimes dt + \sum_{i=3}^n \left(\prod_{s \in S} H_s^{-2h_s \delta_{iI_s}} \right) g^i \right\},$$

$$\exp(\varphi^\alpha) = \prod_{s \in S} H_s^{h_s \chi_s \lambda_{a_s}^\alpha}, \quad (5)$$

$$F^a = \sum_{s \in S} \delta_{a_s}^a \mathcal{F}^s, \quad (6)$$

where $f = 1 - 2\mu/R^{\bar{d}}$,

$$\mathcal{F}^s = -\frac{Q_s}{R^{d_1}} \left(\prod_{s' \in S} H_{s'}^{-A_{ss'}} \right) dR \wedge \tau(I_s), \quad s \in S_e, \quad (7)$$

$$\mathcal{F}^s = Q_s \tau(\bar{I}_s), \quad s \in S_m. \quad (8)$$

Here $Q_s \neq 0$ ($s \in S$) are charges, $R_0 > 0$, $R_0^{\bar{d}} = 2\mu > 0$ and $\bar{d} = d_1 - 1$. In (4) $g^i = g_{m_i n_i}^i(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$ is a Ricci-flat metric on M_i , $i = 3, \dots, n$ and $\delta_{iI} = \sum_{j \in I} \delta_{ij}$ is the indicator of i belonging to I : $\delta_{iI} = 1$ for $i \in I$ and $\delta_{iI} = 0$ otherwise. Here $g^2 = -dt \otimes dt$, and $g^1 = d\Omega_{d_1}$ is a canonical metric on unit sphere $M_1 = S^{d_1}$,

The p -brane set S is by definition

$$S = S_e \cup S_m, \quad S_v = \cup_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (9)$$

$v = e, m$ and $\Omega_{a,e}, \Omega_{a,m} \subset \Omega$, where $\Omega = \Omega(n)$ is the set of all non-empty subsets of $\{2, \dots, n\}$, i.e. all p -branes do not “live” in M_1 .

Any p -brane index $s \in S$ has the form $s = (a_s, v_s, I_s)$, where $a_s \in \Delta$, $v_s = e, m$ and $I_s \in \Omega_{a_s, v_s}$. The sets S_e and S_m define electric and magnetic p -branes, correspondingly. In (5) $\chi_s = +1, -1$ for $s \in S_e, S_m$, respectively. All p -branes contain the time manifold $M_2 = \mathbb{R}$, i.e.

$$2 \in I_s, \quad \forall s \in S. \quad (10)$$

All manifolds M_i , $i > 2$, are oriented and connected and

$$\tau_i \equiv \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad (11)$$

are volume d_i -forms, where $d_i = \dim M_i$, $i = 1, \dots, n$, with $d_1 > 1$ and $d_2 = 1$. For any $I = \{i_1, \dots, i_k\} \in \Omega$, $i_1 < \dots < i_k$, we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k}, \quad d(I) = \sum_{i \in I} d_i. \quad (12)$$

The forms $\mathcal{F}^s$ correspond to the electric and magnetic p -branes for $s \in S_e, S_m$, respectively. In (8) we use the notation $\bar{I} = \{1, \dots, n\} \setminus I$.

The parameters h_s appearing in the solution satisfy the relations: $h_s = (B_{ss})^{-1}$, where

$$B_{ss'} = d(I_s \cap I_{s'}) + \frac{d(I_s)d(I_{s'})}{2-D} + \chi_s \chi_{s'} \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta}, \quad (13)$$

$s, s' \in S$, with $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$ and $D = 1 + \sum_{i=1}^n d_i$. Here we assume that

$$(i) \quad B_{ss} \neq 0,$$

$s \in S$, and

$$(ii) \quad \det(B_{ss'}) \neq 0,$$

i.e. the matrix $(B_{ss'})$ is a non-degenerate one.

Let us consider the matrix

$$(A_{ss'}) = (2B_{ss'}/B_{s's'}). \quad (14)$$

Here some ordering in S is assumed.

The functions $H_s = H_s(z) > 0$, $z = 2\mu/R^{\bar{d}} \in (0, 1)$, obey the equations

$$\frac{d}{dz} \left(\frac{(1-z)}{H_s} \frac{dH_s}{dz} \right) = B_s \prod_{s' \in S} H_{s'}^{-A_{ss'}}, \quad (15)$$

equipped with the boundary conditions

$$H_s(1-0) = H_{s0} \in (0, +\infty), \quad (16)$$

$$H_s(+0) = 1, \quad (17)$$

$s \in S$. Here

$$B_s = B_{ss} \varepsilon_s Q_s^2 / (2\bar{d}\mu)^2, \quad \varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \quad (18)$$

$s \in S$, $\varepsilon[g] \equiv \text{sign det}(g_{MN})$. More explicitly eq. (18) reads: $\varepsilon_s = \varepsilon(I_s) \theta_{a_s}$ for $v_s = e$ and $\varepsilon_s = -\varepsilon[g] \varepsilon(I_s) \theta_{a_s}$, for $v_s = m$.

Eqs. (15) had to be solved in order to obtain explicit solutions. They are equivalent to the Toda-type equations [4]. The first boundary condition (16) guarantees the existence of a regular horizon at $R^{\bar{d}} = 2\mu$. The second condition (17) ensures an asymptotical flatness (for $R \rightarrow +\infty$) of the $(2 + d_1)$ -dimensional section of the metric.

Due to eqs. (7) and (8), the dimension of p -brane world volume $d(I_s)$ is defined by relations: $d(I_s) = n_{a_s} - 1$, $d(I_s) = D - n_{a_s} - 1$, for $s \in S_e, S_m$, respectively. For a p -brane we use a standard notation: $p = p_s = d(I_s) - 1$.

The solutions are valid if the appropriate restriction on the sets $\Omega_{a,v}$ is imposed. This restriction guarantees the block-diagonal structure of the stress-energy tensor. We denote $w_1 \equiv \{i | i \in \{2, \dots, n\}, d_i = 1\}$ and $n_1 = |w_1|$ (i.e. n_1 is the number of 1-dimensional spaces among M_i , $i = 2, \dots, n$). It is clear, that $2 \in w_1$.

Restriction. Let 1a) $n_1 \leq 1$ or 1b) $n_1 \geq 2$ and for any $a \in \Delta$, $v \in \{e, m\}$, $i, j \in w_1$, $i \neq j$, there are no $I, J \in \Omega_{a,v}$ such that $i \in I$, $j \in J$ and $I \setminus \{i\} = J \setminus \{j\}$.

This restriction is satisfied in the non-composite case: $|\Omega_{a,e}| + |\Omega_{a,m}| = 1$, (i.e when there are no two p -branes with the same color index a , $a \in \Delta$.) The restriction forbids certain intersections of two p -branes with the same color index for $n_1 \geq 2$.

The Hawking temperature corresponding to the solution has the form [1]

$$T_H = \frac{\bar{d}}{4\pi(2\mu)^{1/\bar{d}}} \prod_{s \in S} H_{s0}^{-h_s}, \quad (19)$$

where H_{s0} are defined in (16).

The solution under consideration describes a set of charged (by forms) overlapping black p -branes "living" on submanifolds of $M_2 \times \dots \times M_n$.

3 Examples of solutions

3.1 "Block-orthogonal" solutions

The simplest polynomial solutions of eqs. (15) occur in orthogonal case [8, 9, 10, 12, 13], when

$$B_{ss'} = 0, \quad (20)$$

for $s \neq s'$, $s, s' \in S$. In this case $(A_{ss'}) = \text{diag}(2, \dots, 2)$ is a Cartan matrix for semisimple Lie algebra $A_1 \oplus \dots \oplus A_1$ and

$$H_s(z) = 1 + P_s z, \quad (21)$$

with $P_s \neq 0$, satisfying

$$P_s(P_s + 1) = -B_s, \quad (22)$$

$s \in S$. For positive parameters $P_s > 0$ we get negative $B_s < 0$.

In [15, 16] this solution of eqs. (15) was generalised to the "block-orthogonal" case:

$$S = S_1 \cup \dots \cup S_k, \quad S_i \cap S_j = \emptyset, \quad i \neq j, \quad (23)$$

$S_i \neq \emptyset$, i.e. the set S is a union of k non-intersecting (non-empty) subsets $S_1, \dots, S_k$, and relation (20) is satisfied for all $s \in S_i$, $s' \in S_j$, $i \neq j$; $i, j = 1, \dots, k$. In this case eq. (21) is modified as follows

$$H_s(z) = (1 + P_s z)^{b^s}, \quad (24)$$

where

$$b^s = 2 \sum_{s' \in S} A^{ss'}, \quad (25)$$

$(A^{ss'}) = (A_{ss'})^{-1}$ and parameters P_s are coinciding inside blocks, i.e. $P_s = P_{s'}$ for $s, s' \in S_i$, $i = 1, \dots, k$. Parameters $P_s \neq 0$ satisfy the relations (22) and parameters B_s are also coinciding inside blocks, i.e. $B_s = B_{s'}$ for $s, s' \in S_i$, $i = 1, \dots, k$.

Let $(A_{ss'})$ be a Cartan matrix for a finite-dimensional semisimple Lie algebra $\mathcal{G}$. In this case all the powers in (25) are natural numbers coinciding with the components of twice the dual Weyl vector in the basis of simple roots [17], and hence, all functions H_s are polynomials, $s \in S$. The following conjecture may help in finding solutions of eqs. (15).

Conjecture [1]. *Let $(A_{ss'})$ be a Cartan matrix for a semisimple finite-dimensional Lie algebra $\mathcal{G}$. Then the solution to eqs. (15)-(17) (if exists) is a polynomial*

$$H_s(z) = 1 + \sum_{k=1}^{n_s} P_s^{(k)} z^k, \quad (26)$$

where $P_s^{(k)}$ are constants, $k = 1, \dots, n_s$, the integers $n_s = b^s$ are defined in (25) and $P_s^{(n_s)} \neq 0$, $s \in S$.

This conjecture was verified for A_n and C_n series of Lie algebras [2, 3]. In the extremal case $\mu = +0$ an analogue of this conjecture was suggested earlier in [11].

3.2 Solution for A_2 algebra

Here before coming to solutions for A_3 algebra we present the polynomial solution from [1, 2] corresponding to the Lie algebra $A_2 = sl(3)$ with the Cartan matrix

$$(A_{ss'}) = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}. \quad (27)$$

The moduli polynomials read in this case as follows

$$H_s = 1 + P_s z + P_s^{(2)} z^2, \quad (28)$$

where $P_s = P_s^{(1)}$ and $P_s^{(2)} \neq 0$ are constants and

$$P_s^{(2)} = \frac{P_s P_{s+1} (P_s + 1)}{2(P_1 + P_2 + 2)}, \quad (29)$$

$$B_s = -\frac{P_s (P_s + 1) (P_s + 2)}{P_1 + P_2 + 2}, \quad (30)$$

$s = 1, 2$. Here $P_1 + P_2 + 2 \neq 0$.

In the A_2 -case the solution is described by relations (4)-(8) with $S = \{s_1, s_2\}$ and intersection rules are following from (13), (14) and (27)

$$d(I_{s_1} \cap I_{s_2}) = \frac{d(I_{s_1})d(I_{s_2})}{D-2} - \chi_{s_1} \chi_{s_2} \lambda_{a_{s_1}} \cdot \lambda_{a_{s_2}} - \frac{1}{2}K, \quad (31)$$

$$d(I_{s_i}) - \frac{(d(I_{s_i}))^2}{D-2} + \lambda_{a_{s_i}} \cdot \lambda_{a_{s_i}} = K, \quad (32)$$

where $K \neq 0$ and functions $H_{s_i} = H_i$ are defined by relations (28)-(30) with $z = 2\mu R^{-\bar{d}}$, $i = 1, 2$. Here and in what follows $\lambda \cdot \lambda' = \lambda_\alpha \lambda'_\beta h^{\alpha\beta}$.

3.3 Solution for A_3 algebra

Now we present the new solution related to the Lie algebra $A_3 = sl(4)$ with the Cartan matrix

$$(A_{ss'}) = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}. \quad (33)$$

According to **Conjecture** we seek the solution to eqs. (15)-(17) in the following form

$$H_1(z) = 1 + P_1 z + P_1^{(2)} z^2 + P_1^{(3)} z^3, \quad (34)$$

$$H_2(z) = 1 + P_2 z + P_2^{(2)} z^2 + P_2^{(3)} z^3 + P_2^{(4)} z^4, \quad (35)$$

$$H_3(z) = 1 + P_3 z + P_3^{(2)} z^2 + P_3^{(3)} z^3, \quad (36)$$

where $P_s = P_s^{(1)}$ and $P_s^{(k)}$ are constants, $s = 1, 2, 3$.

Here we outline the result obtained by computer calculations. For B_s -parameters (18) we get the following relations

$$B_1 = \frac{1}{2(2 + P_1 + P_2 - P_3)} \left[12 - \Delta - 2P_1^3 - 2P_1^2(3 - P_3) + 6P_3 \right. \\ \left. + P_2(4 + P_2)(3 + P_3) + P_1(2 + P_2(4 + P_2) + 6P_3) \right], \quad (37)$$

$$B_3 = \frac{1}{2(2 - P_1 + P_2 + P_3)} \left[12 - \Delta + 6P_1 + (3 + P_1)P_2(4 + P_2), \right. \\ \left. + (2 + 6P_1 + P_2(4 + P_2))P_3 - 2(3 - P_1)P_3^2 - 2P_3^3 \right], \quad (38)$$

$$B_2 = \frac{1}{(2 + P_1 + P_2 - P_3)(2 - P_1 + P_2 + P_3)} \left[(2 + P_2)(\Delta \right. \\ \left. - (2 + P_2)(2 + (2 + P_2)^2 + 2P_1P_3 + 3(P_1 + P_3))) \right], \quad (39)$$

and for parameters $P_s^{(k)}$ in (15)-(17) we obtain

$$P_1^{(2)} = \frac{1}{4(2 + P_1 + P_2 - P_3)} \left[12 - \Delta + 2P_1^2P_2 + 6P_3 + P_2(4 + P_2)(3 + P_3) \right. \\ \left. + P_1(6 + P_2(6 + P_2) + 4P_3) \right], \quad (40)$$

$$P_3^{(2)} = \frac{1}{4(2 - P_1 + P_2 + P_3)} \left[12 - \Delta + 6P_1 + (3 + P_1)P_2(4 + P_2) \right. \\ \left. + (6 + 4P_1 + P_2(6 + P_2))P_3 + 2P_2P_3^2 \right], \quad (41)$$

$$P_1^{(3)} = \frac{1}{12(2 + P_1 + P_2 - P_3)} \left[(2 + P_1 + P_2)(12 - \Delta + 6P_1 + (3 + P_1)P_2(4 + P_2)) \right. \\ \left. + (2P_1^2(2 + P_2) + (2 + P_2)(6 + P_2(4 + P_2)) + P_1(14 + P_2(10 + P_2)))P_3 \right], \quad (42)$$

$$P_3^{(3)} = \frac{1}{12(2 - P_1 + P_2 + P_3)} \left[P_1((2 + P_2)(6 + P_2(4 + P_2)) + (14 + P_2(10 + P_2))P_3 \right. \\ \left. + 2(2 + P_2)P_3^2) + (2 + P_2 + P_3)(12 - \Delta + 6P_3 + P_2(4 + P_2)(3 + P_3)) \right], \quad (43)$$

$$P_2^{(2)} = \frac{1}{2(2 + P_1 + P_2 - P_3)(2 - P_1 + P_2 + P_3)} \left[-((2 + P_2)(12 - \Delta + 3P_2(4 + P_2)) \right. \\ \left. + 2P_1(4 + 3P_2)P_3 + 3(2 + P_2)^2(P_1 + P_3) + P_2(1 + P_2)(P_1^2 + P_3^2)) \right], \quad (44)$$

$$P_2^{(3)} = \frac{1}{6(2 + P_1 + P_2 - P_3)(2 - P_1 + P_2 + P_3)} \left[(2 + P_2)(-(P_2^3(3 + P_1 + P_3)) \right. \\ \left. + \Delta(2 + P_1 + P_2 + P_3) - P_2^2(18 + P_1(9 + P_1) + P_3(9 + P_3)) - 2(2 + P_1 + P_3) \right. \\ \left. \times (6 + 2P_1P_3 + 3(P_1 + P_3) + P_2(9 + P_1P_3 + 2(P_1 + P_3)))) \right], \quad (45)$$

$$P_2^{(4)} = \frac{1}{24(2 + P_1 + P_2 - P_3)(2 - P_1 + P_2 + P_3)} \left[(2 + P_2)(-((3 + P_2)(2 + P_2 + P_3) \right. \\ \left. \times (12 - \Delta + 6P_3 + P_2(4 + P_2)(3 + P_3))) + P_1((-2 - P_2)(3 + P_2)^2(4 + P_2) \right. \\ \left. - 2(42 + P_2(42 + P_2(12 + P_2)))P_3 - 2(2 + P_2)(6 + P_2)P_3^2 + \Delta(3 + P_2 + 2P_3)) \right. \\ \left. - P_1^2(P_2^3 + 2(3 + 2P_3)^2 + P_2^2(7 + 2P_3) + P_2(18 + 4P_3(4 + P_3))) \right], \quad (46)$$

where

$$\begin{aligned} \Delta = & \left[P_1^2 \left((6 + P_2(4 + P_2))^2 + 12(2 + P_2)^2 P_3 + 4(2 + P_2)^2 P_3^2 \right) \right. \\ & + 2P_1 \left(3(2 + P_2)^2 (6 + P_2(4 + P_2)) + (84 + P_2(4 + P_2)(26 + P_2(4 + P_2))) P_3 \right. \\ & \left. \left. + 6(2 + P_2)^2 P_3^2 \right) + (6(2 + P_3) + P_2(4 + P_2)(3 + P_3))^2 \right]^{1/2}. \end{aligned} \quad (47)$$

Here $2 + P_2 \pm (P_1 - P_3) \neq 0$.

The A_3 black-brane solution is described by relations (4)-(8), with $S = \{s_1, s_2, s_3\}$, and intersection rules following from (13), (14) and (33)

$$d(I_{s_1} \cap I_{s_2}) = \frac{d(I_{s_1})d(I_{s_2})}{D-2} - \chi_{s_1}\chi_{s_2}\lambda_{a_{s_1}} \cdot \lambda_{a_{s_2}} - \frac{1}{2}K, \quad (48)$$

$$d(I_{s_2} \cap I_{s_3}) = \frac{d(I_{s_2})d(I_{s_3})}{D-2} - \chi_{s_2}\chi_{s_3}\lambda_{a_{s_2}} \cdot \lambda_{a_{s_3}} - \frac{1}{2}K, \quad (49)$$

$$d(I_{s_1} \cap I_{s_3}) = \frac{d(I_{s_1})d(I_{s_3})}{D-2} - \chi_{s_1}\chi_{s_3}\lambda_{a_{s_1}} \cdot \lambda_{a_{s_3}}, \quad (50)$$

$$d(I_{s_i}) - \frac{(d(I_{s_i}))^2}{D-2} + \lambda_{a_{s_i}} \cdot \lambda_{a_{s_i}} = K, \quad (51)$$

where $K \neq 0$ and functions $H_{s_i} = H_i$ are defined by relations (34)-(36) with $z = 2\mu R^{-\bar{d}}$, $i = 1, 2, 3$.

For $P_1 = 3P$, $P_2 = 4P$ and $P_3 = 3P$ with $P > 0$ we get a special “block-orthogonal” solution

$$H_1(z) = (1 + Pz)^3, \quad H_2(z) = (1 + Pz)^4, \quad H_3(z) = (1 + Pz)^3, \quad (52)$$

in agreement with the relations (24) and (25).

In the special case $P_1 = P_3$ the relations for H_1, H_2 (34)-(35) describe the solution corresponding to Lie algebra $C_2 = so(5)$ [18].

3.3.1 Solution with three 3-branes in $D = 12$ model

Now we illustrate the general A_3 formulae by considering a bosonic field model in dimension $D = 12$ [19] that admits the bosonic sector of 11-dimensional supergravity as a consistent truncation. The action for this model has the following form

$$\begin{aligned} S_{12} = & \int_M d^{12}z \sqrt{|g|} \{ R[g] - g^{MN} \partial_M \varphi \partial_N \varphi - \frac{1}{4!} \exp(2\lambda_4 \varphi) (F^4)^2 - \frac{1}{5!} \exp(2\lambda_5 \varphi) (F^5)^2 \} \\ & + c_{12} \int_M A^4 \wedge F^4 \wedge F^4. \end{aligned} \quad (53)$$

Here $F^4 = dA_3$ is the 4-form, $F^5 = dA^4$ is the 5-form, c_{12} is constant and

$$\lambda_4^2 = -\frac{1}{10}, \quad \lambda_5 = -2\lambda_4. \quad (54)$$

The model describes electrically charged 2- and 3-branes and magnetically charged 6- and 5-branes (corresponding to F^4 and F^5 , respectively) with (orthogonal) intersection rules given in [7].

First, we consider the A_3 -solution for the truncated model (i.e. without Chern-Simons term) defined on the manifold

$$M = (2\mu, +\infty) \times (M_1 = S^2) \times (M_2 = \mathbb{R}) \times (M_3 = \mathbb{R}^2) \times (M_4 = \mathbb{R}^3) \times (M_5 = \mathbb{R}^2) \times (M_6 = \mathbb{R}), \quad (55)$$

with three electric 3-branes with sets $I_i = I_{s_i}$ defined as follows

$$I_1 = \{2, 3, 6\}, \quad I_2 = \{2, 4\}, \quad I_3 = \{2, 5, 6\}, \quad (56)$$

The A_3 -intersection rules (48)-(51) read

$$d(I_1 \cap I_2) = d(I_2 \cap I_3) = 1, \quad d(I_1 \cap I_3) = 2, \quad (57)$$

(here $K = 2$). The black-brane solution has the following form

$$g = (H_1 H_2 H_3)^{4/10} \left\{ f^{-1} dR \otimes dR + R^2 d\Omega_2^2 \right. \quad (58)$$

$$\left. - (H_1 H_2 H_3)^{-1} f dt \otimes dt + H_1^{-1} g^3 + H_2^{-1} g^4 + H_3^{-1} g^5 + H_1^{-1} H_3^{-1} g^6 \right\},$$

$$\exp(\varphi) = (H_1 H_2 H_3)^{\lambda_5/2}, \quad (59)$$

$$F^5 = \frac{Q_1}{R^2} H_1^{-2} H_2 dt \wedge dR \wedge \tau_3 \wedge \tau_6 + \frac{Q_2}{R^2} H_1 H_2^{-2} H_3 dt \wedge dR \wedge \tau_4 + \frac{Q_3}{R^2} H_2 H_3^{-2} dt \wedge dR \wedge \tau_5 \wedge \tau_6, \quad (60)$$

and $F^4 = 0$, where $f = 1 - 2\mu/R$.

Here $Q_i \neq 0$ are charges related to parameters P_i , by formulae $B_i = -Q_i^2/(2\mu^2)$, and (37)-(39); $\mu > 0$, and the functions $H_i = H_i(z, (P_j))$ are defined by (34)-(36), (40)-(47), and $z = 2\mu/R$, $i = 1, 2, 3$. In (58) g^a is a flat metric on M_a , $a = 3, \dots, 6$.

This solution satisfies the equations of motion for the model (53) itself, since the only modifications due to Chern-Simons term are related to ‘‘Maxwells’’ equations

$$d * F^4 = \text{const } F^5 \wedge F^4, \quad d * F^5 = \text{const } F^4 \wedge F^4, \quad (61)$$

and are trivial (since $F^4 = 0$).

We remind that the twelve-dimensional model from [19] can be consistently truncated to a ten-dimensional Lagrangian that contains all the *BPS* p -brane solutions of the type *IIB*-theory.

4 Conclusions

In this paper we presented a new black-brane solution with three branes and intersection rules corresponding to the Lie algebra A_3 . The solution is governed by polynomials of degree 3, 4, 3 and the coefficients of polynomials exhibit a special dependence upon the parameters P_1 , P_2 and P_3 due to appearance of the term Δ (see (47)). (The case $P_1 = P_3$, corresponding to Lie algebra $C_2 = so(5)$ was considered recently in [18].)

The A_3 -solution is more complicated then the A_2 -one, which has a rather simple analytical structure. This means that the polynomial solutions for other A_n -algebras ($n > 3$) may be even more complicated.

Other topics of interest is the application of this result to black-brane thermodynamics (e.g. relations for the entropy, Cardy-Verlinde type formulas etc) and analysis of post-Newtonian effects [2]. On this way one may expect to clarify the appearance of the function Δ in the solution. Another problem is to find the polynomial solutions for other Lie algebras (at least for A_k ones).

Here we also considered as an example the A_3 -solution with three 3-branes in 12-dimensional model from [19] (this model imitates the low-energy limit of hypothetical F theory [20]). This solution may be also generalized to the case of B_D -models [13] in dimensions $D \geq 12$.

Acknowledgments

This work was supported in part by the Russian Ministry of Industry, Science and Technology, Russian Foundation for Basic Research (RFFI-01-02-17312-a) and DFG project (436 RUS 113/678/0-1(R)). The results of the work were reported at the Fifth International Conference on Gravitation and Astrophysics of Asian-Pacific countries.

References

- [1] V.D. Ivashchuk and V.N. Melnikov, P-brane black holes for general intersections, *Grav. Cosmol.*, **5**, 4(20) 313-318 (1999); gr-qc/0002085.
- [2] V.D. Ivashchuk and V.N. Melnikov, Black hole p-brane solutions for general intersection rules, *Grav. Cosmol.*, **6**, 1(21), 27-40 (2000); hep-th/9910041.
- [3] V.D. Ivashchuk and V.N. Melnikov, Toda p-brane black holes and polynomials related to Lie algebras. *Class. Quantum Grav.* **17**, 2073-2092 (2000); math-ph/0002048.
- [4] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.*, **18** (2001) R87-R152; hep-th/0110274.
- [5] I.Ya. Aref'eva and O.A. Rytchkov, Incidence matrix description of intersecting p-brane solutions, *Am. Math. Soc. Transl.* **201** 19-38 (2000); hep-th/9612236.
- [6] I.Ya. Aref'eva, M.G. Ivanov and O.A. Rytchkov, Properties of intersecting p-branes in various dimensions, hep-th/9702077.
- [7] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the generalized composite p-branes, *Class. Quantum Grav.* **14**, 3001-3029 (1997); *Corrigenda* **15** (12), 3941 (1998); hep-th/9705036.
- [8] M. Cvetič and A. Tseytlin, Non-extreme black holes from non-extreme intersecting M -branes, *Nucl. Phys. B* **478**, 181-198 (1996); hep-th/9606033.
- [9] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, Non-extremal intersecting p-branes in various dimensions, *Phys. Lett. B* **406**, 44-48 (1997); hep-th/9702079.
- [10] N. Ohta, Intersection rules for non-extreme p-branes, *Phys. Lett. B* **403**, 218-224 (1997); hep-th/9702164.

- [11] H. Lü, J. Maharana, S. Mukherji and C.N. Pope, Cosmological solutions, p-branes and the Wheeler-DeWitt equation, *Phys. Rev. D* **57**, 2219-2229 (1998); hep-th/9707182.
- [12] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnikov, The Reissner-Nordström problem for intersecting electric and magnetic p-branes, *Grav. Cosmol.* **3**, 3(11), 203-212 (1997); gr-qc/9710054.
- [13] V.D. Ivashchuk and V.N. Melnikov, Multidimensional classical and quantum cosmology with intersecting p-branes, *J. Math. Phys.*, **39**, 2866-2889 (1998); hep-th/9708157.
- [14] V.D. Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.*, **41**, 444-460 (2000); hep-th/9907019.
- [15] K.A. Bronnikov, Block-orthogonal brane systems, black holes and wormholes, *Grav. Cosmol.* **4**, 1(13), 49-56 (1998); hep-th/9710207.
- [16] V.D. Ivashchuk and V.N. Melnikov. Multidimensional cosmological and spherically symmetric solutions with intersecting p-branes. In Lecture Notes in Physics, Vol. 537, "Mathematical and Quantum Aspects of Relativity and Cosmology Proceedings of the Second Samos Meeting on Cosmology, Geometry and Relativity held at Pythagoreon, Samos, Greece, 1998, eds: S. Cotsakis, G.W. Gibbons., Berlin, Springer, 2000; gr-qc/9901001.
- [17] J. Fuchs and C. Schweigert, Symmetries, Lie algebras and representations. A graduate course for physicists. Cambridge Univ. Press, Cambridge, 1997.
- [18] M.A. Grebeniuk, V.D. Ivashchuk and S.-W. Kim, Black-brane solutions for C_2 algebra, submitted to J. Math. Phys., hep-th/0111219.
- [19] N. Khviengia, Z. Khviengia, H. Lü, C.N. Pope, Towards a field theory of F-theory, hep-th/9703012; *Class. Quantum Grav.* **15**, 759-773 (1998).
- [20] C. Vafa, Evidence for F-theory, hep-th/9602022; *Nucl. Phys. B* **469**, 403 (1996).